

# Product datasheet for AP06089PU-M

## **DP1 (TFDP1) Rabbit Polyclonal Antibody**

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies	
Applications:	ELISA, IF, IHC, WB	
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on Paraffin sections: 1/50-1/200. Immunofluorescence: 1/50-1/200.	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Specificity:	This antibody detects endogenous levels of DP-1/TFDP1 protein (region surrounding Val393).	
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% Sodium azide	
Concentration:	1.0 mg/ml	
Purification:	Affinity chromatography using epitope-specific immunogen (> 95% pure by SDS-PAGE)	
Conjugation:	Unconjugated	
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.	
Stability:	Shelf life: one year from despatch.	
Predicted Protein Size:	~55 kDa	
Gene Name:	transcription factor Dp-1	
Database Link:	<u>Entrez Gene 7027 Human</u> <u>Q14186</u>	




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **DP1 (TFDP1) Rabbit Polyclonal Antibody – AP06089PU-M**


Background:	The human retinoblastoma gene product appears to play an important role in the negative egulation of cell proliferation. Functional inactivation of Rb can be mediated either through mutation or as a consequence of interaction with DNA tumor virus-encoded proteins. Of all the Rb associations described to date, the identification of a complex between Rb and the transcription factor E2F most directly implicates Rb in regulation of cell proliferation. E2F was originally identified through its role in transcriptional activation of the adenovirus E2 promoter. Sequences homologous to the E2F binding site have been found upstream of a number of genes that encode proteins with putative functions in the G1 and S phases of the cell cycle. E2F-1 forms heterodimers with a second protein, designated DP-1, forming an "active" E2F transcriptional regulatory complex.
Synonyms:	Transcription factor Dp 1 DDTE1

Synonyms:	Transcription factor Dp-1, DP1, DRTF1
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Cell cycle, TGF-beta signaling pathway

### **Product images:**



Western blot (WB) analysis of TFDP1 antibody (Cat.-No.: [AP06089PU-N]) at 1/500 dilution Lane 1:HEK293T cell lysate Lane 2:Mouse heart tissue lysate Lane 3:PC12 cell lysate

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemical analysis in paraffinembedded human breast carcinoma tissue using TFDP1 antibody (Cat.-No. [AP06089PU-N]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US