

# Product datasheet for AP06121PU-M

## FHIT Rabbit Polyclonal Antibody

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type:           | Primary Antibodies   |
|-------------------------|--|
| Applications:           | IF, IHC, WB  |
| Recommended Dilution:   | Western blot: 1/500-1/1000.<br>Immunohistochemistry on paraffin sections: 1/50-1/200.<br>Immunofluorescence: 1/50-1/200.   |
| Reactivity:             | Human  |
| Host:                   | Rabbit   |
| Clonality:              | Polyclonal   |
| Immunogen:              | Synthetic peptide, corresponding to amino acids 100-150 of Human FHIT.   |
| Specificity:            | This antibody detects endogenous levels of FHIT protein.<br>(region surrounding Glu116)  |
| Formulation:            | Phosphate buffered saline (PBS), pH 7.2.<br>State: Aff - Purified<br>State: Liquid purified lg fraction (> 95% pure by SDS-PAGE)<br>Preservative: 0.05% Sodium azide |
| Concentration:          | 1.0 mg/ml  |
| Purification:           | Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-<br>PAGE)   |
| Conjugation:            | Unconjugated   |
| Storage:                | Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.<br>Avoid repeated freezing and thawing.   |
| Stability:              | Shelf life: one year from despatch.  |
| Predicted Protein Size: | ~ 17 kDa   |
| Gene Name:              | fragile histidine triad  |
| Database Link:          | <u>Entrez Gene 2272 Human</u><br><u>P49789</u>   |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


#### **GRIGENE** FHIT Rabbit Polyclonal Antibody – AP06121PU-M

#### **Background:** FHIT, a candidate tumor suppressor gene, contains the FRA3B common fragile site and is highly susceptible to carcinogen damage. The pattern of mutational inactivation seen with the FHIT gene is unique compared with other known tumor suppressors. FHIT gene structure and expression have been shown to be altered in esophageal, head, neck, lung, gastric, breast, and cervical carcinomas. It has been demonstrated that FHIT exon loss is associated with smoking duration or asbestos exposure. The FHIT protein is a member of the histidine triad (HIT) superfamily and functions as a dinucleoside 5',5'''-P1,P3-triphosphate hydrolase.


#### Synonyms:

Dinucleosidetriphosphatase, AP3Aase

#### **Product images:**



Western blot (WB) analysis of FHIT antibody in extracts from A549 cells.



Immunohistochemistry (IHC) analyzes of FHIT antibody in paraffin-embedded human breast carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US