

Product datasheet for AP06125PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

c Fos (FOS) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on paraffin sections: 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 1-50 of Human c-Fos.

Specificity: This antibody detects endogenous levels of Fos protein.

(region surrounding Phe3)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE).

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: Fos proto-oncogene, AP-1 transcription factor subunit

Database Link: Entrez Gene 2353 Human

P01100






Background:


The c-Fos oncogene was initially identified as the transforming gene of two independent murine osteosarcoma virus isolates and an avian nephroblastoma virus. The cellular homolog, c-Fos, encodes a nuclear phosphoprotein that is rapidly and transiently induced by a variety of agents and functions as a transcriptional regulator for several genes. In contrast to c-Jun proteins, which form homo- and heterodimers which bind to specific DNA TPA response elements (TREs), c-Fos proteins are only active as heterodimers with members of the Jun gene family. Murine Fos B encodes a nuclear protein of 338 amino acids which has 70% homology with c-Fos, exhibits similar kinetics of expression as c-Fos and forms heterodimers with both c-Jun and Jun B, which bind to TRE DNA response elements. Functional homologs of c-Fos and Fos B include Fra-1 and Fra-2 genes.

Synonyms: FOS, G0S7

Product images:



Western blot (WB) analysis of c-Fos antibody at 1/500 dilution Lane 1: Hela whole cell lysate Lane 2: Mouse brain tissue lysate Lane 3: Mouse herat tissue lysate Lane 5: Rat kidney tissue lysate Lane 6: Rat brain tissue lysate



Immunohistochemistry (IHC) analyzes of Fos antibody in paraffin-embedded human breast carcinoma tissue.