

Product datasheet for AP06136PU-M

GDNF Receptor alpha 1 (GFRA1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies IHC, WB **Applications:** Recommended Dilution: Western blot: 1/500-1/1000. Immunohistochemistry on Paraffin Sections: 1/50-1/200. **Reactivity:** Human, Mouse, Rat Host: Rabbit **Clonality:** Polyclonal Synthetic peptide, corresponding to amino acids 50-100 of Human GFRα-1. Immunogen: This antibody detects endogenous levels of GFR alpha-1 protein. Specificity: (region surrounding Lys90) Formulation: Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE) Preservative: 0.05% Sodium Azide **Concentration:** 1.0 mg/ml **Purification:** Affinity Chromatography using epitope-specific immunogen **Conjugation:** Unconjugated Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage: Avoid repeated freezing and thawing. Stability: Shelf life: one year from despatch. Predicted Protein Size: ~ 51 kDa Gene Name: GDNF family receptor alpha 1 Database Link: Entrez Gene 14585 MouseEntrez Gene 25454 RatEntrez Gene 2674 Human P56159



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


	GDNF Receptor alpha 1 (GFRA1) Rabbit Polyclonal Antibody – AP06136PU-M
Background:	Glial cell line-derived neurotrophic factor (GDNF) and the related neurotrophic factor neurturin (NTN) are potent survival factors for central and peripheral neurons. GDNF is a glycosylated, disulfide-bonded homodimer that is distantly related to the TGF β superfamily of growth factors. Three receptors for these factors, GFR α -1 (also designated GDNFR- α , RETL1 or TrnR-1), GFR α -2 (also designated GDNFR- β , RETL2, NTNR- α or TrnR-2) and GFR α -3 have been identified. The receptors do not contain transmembrane domains and are attached to the cell membrane by glycosyl-phosphoinositol linkage. Both GFR α -1 and GFR α -2 have been shown to mediate the GDNF-dependent and NTN-dependent phosphorylation and activation of the tyrosine kinase Ret. GFR α -3 is expressed only during development.
Synonyms	GER-alpha-1 GDNF receptor alpha RET ligand 1 GDNERA RETU1 TRNR1

Synonyms:GFR-alpha-1, GDNF receptor alpha, RET ligand 1, GDNFRA, RETL1, TRNR1Protein Families:Druggable Genome

Product images:



Western blot (WB) analysis of GFRa-1 antibody at 1/500 dilution Lane 1:HEK293T whole cell lysate Lane 2:Mouse liver tissue lysate Lane 3:PC12 whole cell lysate Lane 4:H9C2 whole cell lysate



Immunohistochemistry analysis of GFRA1 / GDNFR-alpha Antibody in paraffin-embedded human brain tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US