

Product datasheet for AP06141PU-M

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

GRIK2 (841-890) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to amino acids 841-890 of Human mGluR-6.

Specificity: This antibody detects endogenous levels of GluR6 protein.

(region surrounding Ala865)

Formulation: PBS with 0.02% sodium azide, 50% glycerol, pH7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 100 kDa

Gene Name: glutamate ionotropic receptor kainate type subunit 2

Database Link: Entrez Gene 2898 Human

Q13002






Background:

The mGluR proteins (metabotropic glutamate receptors) are members of the G protein-coupled receptor family and are functionally and pharmacologically distinct from the GluR proteins (ionotropic glutamate receptors). The eight currently known mGluR proteins are mediated by two G proteins with opposing regulation of adenylate cyclase pathways. The activities of mGluR-1 and mGluR-5 are mediated by a G protein that activates a phosphatidylinositolcalcium second messenger system and generates a calcium-activated chloride current. The remainder of the eight subtypes of mGluR have an activity mediated by a G protein that inhibits adenylate cyclase activity. mGluR-6 is expressed in the synapses of bipolar cell dendrites. This receptor is involved in mediating synaptic transmission from rod and cone photoreceptors to other neurons.


Synonyms:

GRIK2, Excitatory amino acid receptor 4, EAA4

Product images:




Western blot (WB) analysis of mGluR-6 antibody at 1/500 dilution: Lane 1: Hela whole cell lysate Lane 2: NIH-3T3 whole cell lysate Lane 3: Raw264.7 whole cell lysate Lane 4: H9C2 whole cell lysate



Immunohistochemistry (IHC) analysis of GluR6 antibody in paraffin-embedded human breast carcinoma tissue at 1/50 dilution.





Immunohistochemistry (IHC) analysis of GluR6 antibody in paraffin-embedded human brain tissue.