

Product datasheet for AP06177PU-M

IGF1 Receptor (IGF1R) Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000.
	Immunohistochemistry on paraffin sections 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 1121-1170 of Human IGF-1R.
Specificity:	This antibody detects endogenous levels of IGF-1R protein. (region surrounding lle1160)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 15 mM Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 95,155 kDa
Gene Name:	insulin like growth factor 1 receptor
Database Link:	<u>Entrez Gene 3480 Human</u> <u>P08069</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

IGF1 Receptor (IGF1R) Rabbit Polyclonal Antibody - AP06177PU-MBackground:Insulin-like growth factor-I receptor (IGF-IR) is an RTK that stimulates growth in many different
cell types, blocks apoptosis, acts as an intermediate of many growth hormone responses and
may stimulate the growth of some types of cancer. The IGF-IR cognate ligand insulin-like
growth factor-I (IGF-I) promotes association of IGF-IR with Shc, GRB2 and Sos 1, which initiates
Ras and ERK kinase cascades, thereby modifying transcription factor activity, such as
activation of the Elk transcription factors. The modular phosphotyrosine binding (PTB)
domains of insulin receptor substrate (IRS)-1 and -2 can associate with active IGF-IR and
initiate phosphatidylinositol 3-kinase-dependent downstream signals.


Synonyms: IGF-I receptor, IGF1 Receptor

Product images:



Western blot (WB) analysis of IGF-1R antibody

Lane1:MCF-7 whole cell lysate Lane2:HEK293T whole cell lysate Lane3:Hela whole cell lysate IGF-1R (G1155) pAb at 1:500 dilution



Immunohistochemistry (IHC) analyzes of IGF-1R antibody in paraffin-embedded human breast carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US