

Product datasheet for AP06187PU-M

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

COPS3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunofluorescence: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 376-425 of Human COPS3.

Specificity: This antibody detects endogenous levels of JAB1/COPS3 protein.

(region surrounding Met408)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 47 kDa

Gene Name: COP9 signalosome subunit 3

Database Link: Entrez Gene 8533 Human

O9UNS2






Background:

Genes belonging to the Jun and Fos oncogene families encode nuclear proteins that are found to be associated with a number of transcriptional complexes. The c-Jun protein is a major component of the transcription factor AP-1, originally shown to mediate phorbol ester tumor promoter (TPA)-induced expression of responsive genes through the TPA-response element (TRE). The Jun proteins form homo- and heterodimers which bind the TRE, but the Fos proteins are active only as heterodimers with any of the Jun proteins. Fos/Jun heterodimers have a much higher affinity for the TRE than Jun homodimers. Ha-Ras augments c-Jun activity and stimulates phosphorylation of its activation domain. The coactivator of Jun, designated JAB1 (for Junactivation domain-binding protein), interacts with c-Jun and Jun D, but not with Jun B or v-Jun. This interaction enhances the transactivating ability of Jun proteins by stabilizing their binding to the TRE.

Synonyms:

Signalosome subunit 3, SGN3, CSN3

Product images:



Western blot (WB) analysis of COPS3 antibody at 1/500 dilution Lane 1:Hela whole cell lysate Lane 2:sp2/0 whole cell lysate Lane 3:Mouse liver tissue lysate Lane 4:Rat liver tissue lysate