

Product datasheet for AP06197PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cytokeratin 16 (KRT16) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on Paraffin Sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 421-470 of Human Cytokeratin 16.

Specificity: This antibody detects endogenous levels of Cytokeratin 16 protein.

(region surroundign Ser451)

Formulation: Phosphate buffered saline (PBS), pH 7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)

Stabilizer: 50 % Glyerol

Preservative: 0.02% Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Predicted Protein Size: ~52 kDa

Gene Name: keratin 16

Database Link: Entrez Gene 3868 Human

P08779





Background:


Keratin 16 is expressed in keratinocytes, which are undergoing rapid turnover in the suprabasal region (also known as hyperproliferation-related keratins). Keratin 16 is absent in normal breast tissue and in noninvasive breast carcinomas. Only 10% of the invasive breast carcinomas show diffuse or focal positivity. Reportedly, a relatively high concordance was found between the carcinomas immunostaining with the basal cell and the hyperproliferation-related keratins, but not between these markers and the proliferation marker Ki-67. This supports the conclusion that basal cells in breast cancer may show extensive proliferation, and that absence of Ki-67 staining does not mean that (tumor) cells are not proliferating.

Two cytokeratins, 6 and 16, are expressed when keratincytes are undergoing rapid turnover in various pathological states, wound healing, psoriasis and some carcinomas


Synonyms:

Cytokeratin-16, Keratin-16, Keratin 16, KRT16, KRT16A, CK16, K16

Product images:



Western blot analysis in extracts from HepG2 cells using Cytokeratin 16 antibody.



Immunohistochemistry analysis in paraffinembedded human breast carcinoma tissue using Cytokeratin 16 antibody.