

Product datasheet for AP06224PU-N

MMP1 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. Immunofluorescence: 1/50-1/200.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 411-460 of Human MMP-1.
Specificity:	This antibody detects endogenous levels of MMP-1 protein. (region surrounding His440)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS- PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 54 kDa
Gene Name:	matrix metallopeptidase 1
Database Link:	<u>Entrez Gene 4312 Human</u> <u>P03956</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE MMP1 Rabbit Polyclonal Antibody – AP06224PU-N

Background:The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the
degradation of extracellular matrix components, including collagen, gelatin, fibronectin,
laminin and proteoglycan. Transcription of MMP genes is differentially activated by phorbol
ester, lipopolysaccharide (LPS) or staphylococcal enterotoxin B (SEB). MMP catalysis requires
both calcium and zinc. MMP-9 (also designated 92 kDa type IV collagenase or gelatinase B)
has been shown to degrade bone collagens in concert with MMP-1 (also designated interstitial
collagenase, fibroblast collagenase or collagenase-1), and cysteine proteases and may play a
role in bone osteoclastic resorption. MMP-1 is downregulated by p53, and abnormality of p53
expression may contribute to joint degradation in rheumatoid arthritis by regulating MMP-1
expression.

Synonyms:

Interstitial collagenase, Matrix metalloproteinase-1, Fibroblast collagenase, MMP1, CLG

Product images:



Western blot (WB) analysis of MMP-1 antibody at 1/500 dilution Lane 1:Hela whole cell lysate treated with LPS Lane 2:Raw264.7 whole cell lysate treated with LPS Lane 3:PC12 whole cell lysate treated with LPS

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US