

Product datasheet for **AP06250PU-N**

NF-kB p65 (RELA) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 500-550 of Human NFkB-p65.
Specificity:	This antibody detects endogenous levels of NFkappaB-p65 protein. (region surrounding Leu523)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 60, 75 kDa
Gene Name:	RELA proto-oncogene, NF-kB subunit
Database Link:	Entrez Gene 5970 Human Q04206




[View online »](#)

Background:


Proteins encoded by the v-Rel viral oncogene and its cellular homolog, c-Rel, are members of a family of transcription factors that include the two subunits of the transcription factor NFkappaB (p50 and p65) and the Drosophila maternal morphogen, dorsal. Both proteins specifically bind to DNA sequences that are the same or slight variations of the 10 bp kappaB sequence in the immunoglobulin kappa light chain enhancer. This same sequence is also present in a number of other cellular and viral enhancers. The DNA binding activity of NFkappaB is activated and rapidly transported from the cytoplasm to the nucleus in cells exposed to mitogens or growth factors. cDNAs encoding precursors for two distinct 50 kDa proteins have been described, designated p105 and p100, respectively. The p105 precursor contains p50 at its amino terminus and a C-terminal region that when expressed as a separate molecule, designated pd1, binds to p50 and regulates its activity.

Synonyms:

NF kappa B p65, NFkB p65, Transcription factor p65, Rel A, NFKB3

Product images:

Western blot (WB) analysis of NFkB-p65 antibody at 1/500 dilution Lane 1:DLD whole cell lysate Lane 2:Mouse heart tissue lysate Lane 3:Rat brain tissue lysate



Immunohistochemistry (IHC) analyzes of NFkB-p65 antibody in paraffin-embedded human breast carcinoma tissue.