

# Product datasheet for AP06262PU-S

# p18 INK4c (CDKN2C) Rabbit Polyclonal Antibody

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 100-150 of Human p18 INK4c.
Specificity:	This antibody detects endogenous levels of p18 INK protein. (region surrounding Asp142)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 15 mM sodium azide
Concentration:	1.0 mg/ml
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 18 kDa
Gene Name:	cyclin-dependent kinase inhibitor 2C
Database Link:	<u>Entrez Gene 12580 MouseEntrez Gene 1031 Human</u> <u>P42773</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### Sourigene p18 INK4c (CDKN2C) Rabbit Polyclonal Antibody – AP06262PU-S


- Background:The normal progression of cells through the cell cycle is under the control of the cyclin<br/>dependent protein kinases Cdk4 and Cdk6 which are subject to inhibition by the mitotic<br/>inhibitory protein p16. Isolated members of the p16 family have been designated p15 and<br/>p18, respectively. The p16 related protein, p18, interacts strongly with Cdk6 and to a lesser<br/>extent with Cdk4, but lacks apparent interaction with other Cdks. Recombinant p18 has been<br/>shown to inhibit cyclin D-Cdk6 kinase activity. In contrast to p21/p27 that form ternary<br/>complexes with cyclin-Cdks, only binary complexes of p15, p16 and p18 have been identified<br/>in association with Cdk4 and/or Cdk6.
- Synonyms:p18-INK4c, p18-INK6, p18INK6, CDKN6Protein Families:Druggable Genome
- Protein Pathways: Cell cycle

### **Product images:**



Western blot (WB) analysis of p18 INK4c antibody in extracts from hela cells.

Hela whole cell lysate p18 (D142) pAb at 1:500 dilution



Immunohistochemistry (IHC) analysis of p18 INK antibody in paraffin-embedded human breast carcinoma tissue

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US