

Product datasheet for AP06268PU-S

p53 (TP53) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: Western blot: 1/500-1/1000.

Reactivity: Human Rabbit Host: Clonality: Polyclonal

Synthetic peptide, corresponding to amino acids 1-50 of Human p53. Immunogen:

Specificity: This antibody detects endogenous levels of p53 protein.

(region surrounding Met40)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 15 mM sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~53,43 kDa

Gene Name: tumor protein p53

Database Link: Entrez Gene 7157 Human

P04637



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

p53 is a DNA-binding, oligomerization domain and transcription activation domain-containing tumor suppressor that upregulates growth arrest and apoptosis-related genes in response to stress signals, thereby influencing programmed cell death, cell differentiation and cell cycle control mechanisms. p53 localizes to the nucleus yet can be chaperoned to the cytoplasm by the negative regulator MDM2, an E3 ubiquitin ligase that is upregulated in the presence of active p53, where MDM2 polyubiquitinates p53 for proteasome targeting. p53 can assemble into tetramers in the absence of DNA, fluctuates between latent and active (DNA-binding) conformations, and is differentially activated through posttranslational modifications including phosphorylation and acetylation. Mutations in the DNA-binding domain (DBD) (amino acids 110-286) of p53 can compromise energetically favorable association with cis elements and are implicated in several human cancers.

Synonyms:

Cellular tumor antigen p53, Tumor suppressor p53, Phosphoprotein p53, NY-CO-13

Product images:



Western blot (WB) analysis of p53 antibody in extracts from THP1 cells.