

### Product datasheet for AP06279PU-S

# **PAX5 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: Western blot: 1/500-1/1000.

Reactivity: Human, Mouse

Rabbit Host:

Clonality: Polyclonal

Synthetic peptide, corresponding to the N-terminus of Human Pax-5. Immunogen:

Specificity: This antibody detects endogenous levels of Pax-5 protein.

(region surrounding Asp2)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 15 mM sodium azide

Concentration: 1.0 mg/ml

**Purification:** Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 40 kDa

Gene Name: paired box 5

Database Link: Entrez Gene 5079 Human

Q02548



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

The Pax gene family of nuclear transcription factors is comprised of nine family members that function during embryogenesis to regulate the temporal and position-dependent differentiation of cells. In addition, the family is involved in a variety of signal transduction pathways in the adult organism. Mutations in the Pax family of proteins have been linked to disease and cancer in humans. For example, the human Pax-5 gene encodes a B cell lineage specific protein called B cell specific activator protein or BSAP, which is expressed in pro-B, pre-B and mature B lymphocytes but not in plasma cells. BSAP functions to regulate not only B cell development, but also influences the balance between immunoglobulin secretion and B cell proliferation. Overexpression of BSAP has been implicated in cellular transformation, and in the case of small lyphocytic lymphomas with plasmacytoid differentiation, a t(9;14) (p13;q32) translocation resulting in the deregulation of Pax-5 gene expression has been detected. The gene which encodes Pax-5 maps to human chromosome 9p13.

**Synonyms:** Pax-5, BSAP

## **Product images:**



Western blot (WB) analysis of Pax-5 antibody in extracts from A549 cells.

A549 whole cell lysate Pax-5 (D2) pAb at 1:500 dilution