

Product datasheet for AP06336PU-M

SYK Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 500-550 of Human Syk.
Specificity:	This antibody detects endogenous levels of SYK protein. (region surrounding Leu519)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS- PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 72 kDa
Gene Name:	spleen tyrosine kinase
Database Link:	<u>Entrez Gene 6850 Human</u> <u>P43405</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE SYK Rabbit Polyclonal Antibody – AP06336PU-M

Background:Syk (spleen tyrosine kinase) is a 635 amino acid protein that contains one protein kinase
domain and two SH2 domains. One of several members of the protein kinase superfamily,
Syk functions as a positive effector of B cell antigen receptor (CD79)-stimulated responses,
coupling CD79 with the movement of one calcium ion through one of two phospho-regulated
pathways. Via its ability to influence CD79 activity and to control the movement of calicum
through the cell, Syk plays an important role in a variety of cellular responses, including
differentiation, phagocytosis, proliferation and B cell development. Syk expression is
upregulated in T cell lymphoma, suggesting a possible role for Syk in tumorigenesis. Two
isoforms of Syk, designated short and long, exist due to alternative splicing events.


Synonyms:

Tyrosine-protein kinase SYK, Spleen tyrosine kinase

Product images:



Western blot (WB) analysis of Syk antibody at 1/500 dilution Lane 1:Hela whole cell lysate Lane 2:Raw264.7 whole cell lysate Lane 3:Rat kidney tissue lysate



Immunohistochemistry (IHC) analyzes of SYK antibody in paraffin-embedded human breast carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US