

Product datasheet for AP06351PU-N

Troduct datasirect for Al 6655 if 6 it

TGF beta 2 (TGFB2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500 - 1/1000.

Immunohistochemistry on Paraffin Sections: 1/50 - 1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 360-400 of Human TGFβ2.

Specificity: This antibody detects endogenous levels of TGF beta-2 protein.

Formulation: Phosphate buffered saline (PBS), pH~7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)

Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

Database Link:

Purification: Affinity Chromatography using epitope-specific immunogen

Entrez Gene 7042 Human

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: transforming growth factor beta 2

dene Name.

P61812



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Transforming growth factor betas (TGF β s) were originally discovered due to their ability to promote anchorage-independent growth of rat NRK fibroblasts in the presence of TGF α . It is now realized that TGF β s mediate many cell-cell interactions that occur during embryonic development. Three TGF β s have been identified in mammals. TGF β 1, TGF β 2 and TGF β 3 are each synthesized as precursor proteins that are very similar in that each is cleaved to yield a 112 amino acid polypeptide that remains associated with the latent portion of the molecules. Biologically active TGF β requires dimerization of the monomers (usually homodimers) and release of the latent peptide portion. Overall, the mature region of the TGF β 3 protein has approximately 80% identity to the mature region of both TGF β 1 and TGF β 2. However, the NH2 terminals or precursor regions of their molecules share only 27% sequence identity.


Synonyms:

Transforming growth factor beta-2, G-TSF, Polyergin, Cetermin

Product images:



Western blot analysis of TGFβ2 Antibody in extracts from MDA-MB-435 cells.



Immunohistochemistry analysis of TGFB2 Antibody in paraffin-embedded human lung carcinoma tissue.