

Product datasheet for AP06355PU-M

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

TIMP2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunofluorescence: 1/50-1/200.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 10-60 of Human TIMP-2.

Specificity: This antibody detects endogenous levels of TIMP2 protein.

(region surrounding Ala47)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 24 kDa

Gene Name: TIMP metallopeptidase inhibitor 2

Database Link: Entrez Gene 7077 Human

P16035






Background:

TIMP-1, TIMP-2, TIMP-3 and TIMP-4 (for tissue inhibitor of metalloproteinases-1, -2, -3 and -4) complex with metalloproteinases such as collagenases, vgelatinases and stromelysins, resulting in irreversible inactivation of the metalloproteinase. TIMP-1 was found to be identical to EPA (erythroidpotentiation activity). Parathyroid hormone has been shown to be a regulator of TIMP-2 in osteoblastic cells. TIMP-3 may be involved in regulating trophoblastic invasion of the uterus as well as in regulating remodeling of the extracellular matrix during the folding of epithelia, and in the formation, branching and expansion of epithelial tubes. TIMP-4 is most highly expressed in heart and low levels of TIMP-4 are expressed in liver, brain, lung, thymus and spleen.

Synonyms: CSC-21K, TIMP-2

Product images:



Western blot (WB) analysis of TIMP2 antibody in extracts from A549 cells.