

Product datasheet for AP06359PU-N

Product datasifeet for AP00359PO-I

VAV1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 140-190 of Human VaV1.

Specificity: This antibody detects endogenous levels of VAV1 protein.

(region surrounding Ala168)

Formulation: Phosphate buffered saline (PBS), pH~7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 100 kDa

Gene Name: vav guanine nucleotide exchange factor 1

Database Link: Entrez Gene 7409 Human

P15498



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The Vav gene was originally identified on the basis of its oncogenic activation during the course of gene transfer assays. The major translational product of the Vav proto-oncogene has been identified as a protein containing an array of structural motifs. Contained within its amino terminus are a helix-loop-helix domain and a leucine zipper motif similar to that of Myc family proteins; deletion of this region of p95Vav causes its oncogenic activation. In addition, p95Vav contains an SH2 domain, which could indicate its role as a substrate for tyrosine kinases. Expression of p95Vav is limited exclusively to cells of hematopoietic origin, including those of the erythroid, lymphoid and myeloid lineages. These results suggest that p95Vav may represent a new type of signal transduction molecule involved in the transduction of tyrosine phosphorylation signaling into transcriptional events.

Synonyms: VAV-1

Product images:

Western blot (WB) analysis of VAV1 antibody in extracts from K562 cells treated with UV 15'.



Immunohistochemistry (IHC) analyzes of VAV1 antibody in paraffin-embedded human lung carcinoma breast carcinoma tissue.