

Product datasheet for **AP06362PU-N**

VEGF Receptor 2 (KDR) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 930-970 of Human VEGFR2.
Specificity:	This antibody detects endogenous levels of VEGFR2 protein. (region surrounding Phe945)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 150, 210 kDa
Gene Name:	kinase insert domain receptor
Database Link:	Entrez Gene 3791 Human P35968




[View online »](#)

Background:


Three cell membrane receptor tyrosine kinases, Flt (also designated VEGF-R1), Flk-1 (also designated VEGF-R2) and Flt-4, putatively involved in the growth of endothelial cells, are characterized by the presence of seven immunoglobulinlike sequences in their extracellular domain. These receptors exhibit high degrees of sequence relatedness to each other as well as lesser degrees of relatedness to the class III receptors including CSF-1/Fms, PDGR, SLFR/Kit and Flt-3/Flk-2. Two members of this receptor class, Flt-1 and Flk-1, have been shown to represent high affinity receptors for vascular endothelial growth factors (VEGFs). On the basis of structural similarity to Flt and Flk-1, it has been speculated that Flt-4 might represent a third receptor for either VEGF or a VEGF-related ligand.

Synonyms:

VEGFR2, FLK1, KDR, VEGF Receptor 2

Product images:

Western blot (WB) analysis of VEGFR2 antibody in extracts from SK-OV3 cells.



Immunohistochemistry (IHC) analyzes of VEGFR2/Flk-1 antibody in paraffin-embedded human breast carcinoma brain tissue.