

Product datasheet for **AP06366PU-S**

ACAT1 (ACACA) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western Blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 50-100 of Human ACC α . The immunogen sequence is around H74, total 14 AA.
Specificity:	This antibody detects endogenous levels of ACC1 protein.
Formulation:	PBS, pH 7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 265 kDa
Gene Name:	acetyl-CoA carboxylase alpha
Database Link:	Entrez Gene 31 Human Q13085




[View online »](#)

Background:


Acetyl-CoA carboxylase (ACC) is a complex multifunctional enzyme system which catalyzes the carboxylation of acetyl-CoA to malonyl-CoA, the rate-limiting step in fatty acid synthesis. Exercise diminishes the activity of acetyl-CoA carboxylase in human muscle. ACC α (ACC1) is the rate-limiting enzyme in the biogenesis of long-chain fatty acids, and ACC β (ACC2) may control mitochondrial fatty acid oxidation. These two isoforms of ACC control the amount of fatty acids in the cells. The catalytic function of ACC α is regulated by phosphorylation (inactive) and dephosphorylation (active) of targeted Serine residues and by allosteric transformation by citrate or palmitoyl-CoA, which serve as the short-term regulatory mechanism of the enzyme. The gene encoding ACC α , which maps to human chromosome 17, encodes the 265 kDa α form of ACC, which is the major ACC in lipogenic tissues. The catalytic core of ACC β is homologous to that of ACC α except for an additional peptide of about 150 amino acids at the N-terminus.

Synonyms:

ACACA, ACAC, ACC1, ACCA

Product images:

Western blot (WB) analysis of ACC α antibody at 1/500 dilution Lane 1: Hela whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: NIH-3T3 whole cell lysate Lane 4: PC12 whole cell lysate



Immunohistochemistry (IHC) analyzes of ACC1 antibody in paraffin-embedded human skeletal muscle brain tissue.