

# **Product datasheet for AP06372PU-M**

#### OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **MEF2A Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunofluorescence: 1/50-1/200.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

**Immunogen:** Synthetic peptide, corresponding to amino acids 380-330 of Human MEF-2.

**Specificity:** This antibody detects endogenous levels of MEF2A protein.

(region surrounding Ile402)

Formulation: Phosphate buffered saline (PBS), pH 7.2 with 0.02% Sodium Azide, 50% Glycerol,

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

**Purification:** Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 55 kDa

**Gene Name:** myocyte enhancer factor 2A

Database Link: Entrez Gene 4205 Human

Q02078





#### Background:

The myocyte enhancer factor-2 (MEF-2) family of transcription factors associated with corepessors or co-activators to regulate development and function of T cells, neuronal cells and muscle cells. Four family members arise from alternatively spliced transcripts, termed MEF-2A, -2B, -2C and -2D. These members bind as homo- and heterodimers to the MEF-2 site in the promoter region of affected genes. Differential regulation in the expression of the four transcripts implies functional distinction for each during embryogenesis and development. The process of differentiation from mesodermal precursor cells to myoblasts has led to the discovery of a variety of tissue-specific factors that regulate muscle gene expression. The myogenic basic helix-loop-helix proteins, including MyoD, myogenin, Myf-5 and MRF-4, are one class of identified factors. A second family of DNA binding regulatory proteins is the myocyte-specific enhancer factor-2 (MEF-2) family. Each of these proteins binds to the MEF-2 target DNA sequence present in the regulatory regions of many muscle-specific genes.

Synonyms:


MEF2, Myocyte-specific enhancer factor 2A

### **Product images:**



Western blot (WB) analysis of MEF2A antibody in extracts from HeLa cells.





Immunohistochemistry (IHC) analyzes of MEF2A antibody in paraffin-embedded human breast carcinoma tissue.