

# Product datasheet for AP06384PU-M

### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **DAXX Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human
Host: Rabbit

Clonality: Polyclonal

**Immunogen:** Synthetic peptide, corresponding to amino acids 630-680 of Human Daxx.

**Specificity:** This antibody detects endogenous levels of Daxx protein.

(region surrounding Lys662)

**Formulation:** Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 15 mM sodium azide

**Concentration:** 1.0 mg/ml

**Purification:** Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~ 78, 81 kDa

**Gene Name:** death-domain associated protein

Database Link: Entrez Gene 1616 Human

Q9UER7






#### Background:

Activation of the cell surface receptor FAS by FAS ligand leads to the initiation of apoptosis, a process necessary for the regulation of the immune system and tissue homeostasis. FAS-mediated apoptosis appears to involve a number of divergent and overlapping pathways. Daxx appears to be a central component of a FAS-mediated apoptotic pathway involving the activation of Jun Nterminal kinase (JNK). Although Daxx itself does not contain a death domain, it specifically binds to the death domain of FAS. Overexpression of Daxx activates the JNK pathway and enhances FAS-mediated apoptosis. The Daxx apoptotic pathway acts cooperatively with but is distinct from the FAS-mediated pathway that involves interactions between the death domaincontaining protein FADD and the cysteine protease FLICE. Unlike the FASFADD- FLICE pathway, the Daxx pathway is sensitive to the apoptotic inhibitor protein Bcl-2.


**Synonyms:** BING2, DAP6, EAP1

## **Product images:**



Daxx (K662) pAb at 1:500 dilution

Western blot (WB) analysis of Daxx antibody in extracts from hela cells.



Immunohistochemistry (IHC) analyzes of Daxx antibody in paraffin-embedded human ovary tissue.