

Product datasheet for **AP06420PU-M**

Tyrosine Hydroxylase (TH) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to the N-terminus of Human TH.
Specificity:	This antibody detects endogenous levels of Tyrosine Hydroxylase protein. (region surrounding Gln25)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 58 kDa
Gene Name:	tyrosine hydroxylase
Database Link:	Entrez Gene 7054 Human P07101




[View online »](#)

Background:


The enzyme tyrosine hydroxylase (TH), also designated tyrosine 3-monooxygenase (TY3H), catalyzes the conversion of tyrosine to L-dopa, which is the rate limiting step in the biosynthesis of catecholamines such as dopamine, adrenalin and noradrenalin. TH is thought to play a role in the pathogenesis of Parkinson's disease, which is associated with reduced dopamine levels. Two transcription factor binding sites in the proximal region of the TH gene, the TPA-responsive element (TRE) and the c-AMP responsive element (CRE), have been implicated in the complex regulation of the TH gene. TH is also known to be upregulated by the glia maturation factor (GMF), a Cdc 10/SWI6 motif-containing protein called V-1, and a variety of additional compounds.

Synonyms:

Tyrosine 3-hydroxylase, TYH

Product images:


Western blot (WB) analysis of Tyrosine Hydroxylase antibody in extracts from 293 cells treated with UV 15' or COS7 cells treated with Heat shock.



Immunohistochemistry (IHC) analyzes of Tyrosine Hydroxylase antibody in paraffin-embedded human breast carcinoma tissue.