

# Product datasheet for AP06428PU-S

# FGFR1 Rabbit Polyclonal Antibody

# **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IF, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunofluorescence: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 621-670 of Human FGFR1.
Specificity:	This antibody detects endogenous levels of FGFR1 protein. (region surrounding lle648)
Formulation:	Phosphate Buffered Saline (PBS), pH 7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 15 mM Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Predicted Protein Size:	~ 120, 145 kDa
Gene Name:	fibroblast growth factor receptor 1
Database Link:	<u>Entrez Gene 14182 MouseEntrez Gene 79114 RatEntrez Gene 2260 Human P11362</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **GRIGENE** FGFR1 Rabbit Polyclonal Antibody – AP06428PU-S

Background:	FGFR1 (fibroblast growth factor receptor 1) is a member of the fibroblast growth factor receptor family containing an Ig-like domain and a tyrosine kinase domain. This receptor has multiple isoforms and is a Type I membrane protein. FGFR1 is a widely expressed membrane receptor, with distinct isoforms expressed in specific tissues. FGFR1 binds fibroblast growth factor and induces mitogenesis and cellular differentiation. Defects in FGFR1 result in Pfeiffer syndrome associated with craniosynostosis. FGFR1 can be modified by phosphorylation and can bind basic/acidic fibroblast factor depending on the receptor isoform. FGFR1 has been shown to interact with N-cadherin and NCAM.
Synonyms:	BFGFR, CEK, FGFBR, FLG, FLT2, HBGFR, BFGFR, bFGF-R-1, FLT-2, N-sam, Proto-oncogene c-Fgr
Protein Families:	Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways:Adherens junction, MAPK signaling pathway, Melanoma, Pathways in cancer, Prostate cancer,<br/>Regulation of actin cytoskeleton

## **Product images:**



Western blot (WB) analysis of FGFR1antibody in extracts from RAW264.7 cells.

RAW264.7 whole cell lysate Flg (I648) pAb at 1:500 dilution

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US