

# **Product datasheet for AP06438PU-N**

#### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **RAF1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunofluorescence: 1/50-1/200.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

**Immunogen:** Synthetic peptide, corresponding to amino acids 581-630 of Human Raf-1.

**Specificity:** This antibody detects endogenous levels of Raf1 protein.

(region surrounding Lys615)

**Formulation:** Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

**Concentration:** 1.0 mg/ml

**Purification:** Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 73 kDa

**Gene Name:** Raf-1 proto-oncogene, serine/threonine kinase

**Database Link:** Entrez Gene 5894 Human

P04049






#### Background:


Several serine/threonine protein kinases have been implicated as intermediates in signal transduction pathways. These include ERK/MAP kinases, ribosomal S6 kinase (Rsk) and Raf-1. Raf-1 is a cytoplasmic protein with intrinsic serine/threonine activity. It is broadly expressed in nearly all cell lines tested to date and is the cellular homolog of v-Raf, the product of the transforming gene of the 3611 strain of murine sarcoma virus. The unregulated kinase activity of the v-Raf protein has been associated with transformation and mitogenesis while the activity of Raf-1 is normally suppressed by a regulatory N-terminal domain. Raf-1 is activated in response to activation of a variety of tyrosine kinase receptors as well as in response to pp60v-Src expression. There is accumulating evidence that Ras p21 may play a role in activation of Raf-1 and may even play the role of the messenger from membrane tyrosine kinases to Raf-1.

**Synonyms:** C-RAF, Raf-1, cRaf

## **Product images:**



Western blot (WB) analysis of Raf1 antibody in extracts from HeLa cells treated with UV 5'.



Immunohistochemistry (IHC) analyzes of Raf1 antibody in paraffin-embedded human lung carcinoma tissue.