

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for AP06455PU-N

# E3 SUMO protein ligase PIAS4 (PIAS4) Rabbit Polyclonal Antibody

# **Product data:**

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	This antibody detects endogenous levels of PIAS4 protein. (region surrounding Asp488)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified lg fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS- PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 56 kDa
Gene Name:	protein inhibitor of activated STAT 4
Database Link:	<u>Entrez Gene 51588 Human</u> <u>Q8N2W9</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### SUMO protein ligase PIAS4 (PIAS4) Rabbit Polyclonal Antibody – AP06455PU-N


Background: The IL-6-type family of cytokines, which includes IL-6 as well as a number of similar cytokines and growth factors, plays a significant role in regulating gene activation, proliferation and differentiation. Transcription factors of the Stat family are known to be involved in this signal transduction pathway, undergoing phosphorylation, dimerization and translocation to the nucleus upon activation. PIAS 1, for protein inhibitor of activated Stat1 (also designated Gu/RNA helicase II binding protein), binds specifically to Stat1, blocking Stat1 DNA-binding activity and inhibiting Stat1-mediated gene activation. PIAS 1 also binds to the Gu/RNA helicase II enzyme, leading to the proteolytic cleavage of Gu/RH-II. PIAS 3 similarly binds specifically to Stat3, blocking Stat3 DNA-binding activity and inhibiting Stat3-mediated gene activation.

Synonyms: PIASG, PIAS-gamma, PIASy

### **Product images:**



Western blot (WB) analysis of PIAS4 antibody (Cat.-No.: AP06455PU-N) in extracts from Jurkat cells.



Immunohistochemistry (IHC) analyzes of PIAS4 antibody (Cat.-No.: AP06455PU-N) in paraffinembedded human lung carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US