

## Product datasheet for **AP06477PU-S**

### VEGFB Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	<b>Western blot:</b> 1/500-1/1000. <b>Immunohistochemistry on paraffin sections:</b> 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 100-150 of Human VEGFB.
Specificity:	This antibody detects endogenous levels of VEGFB protein. (region surrounding Arg125)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 22 kDa
Gene Name:	vascular endothelial growth factor B
Database Link:	<a href="#">Entrez Gene 7423 Human P49765</a>




[View online »](#)

**Background:**

The onset of angiogenesis is believed to be an early event in tumorigenesis and may facilitate tumor progression and metastasis. Several growth factors with angiogenic activity have been described. These include fibroblast growth factor (FGF), platelet derived growth factor (PDGF) and vascular endothelial growth factor (VEGF). VEGF is a dimeric glycoprotein with structural homology to PDGF. Several variants of VEGF have been described that arise by alternative mRNA splicing. It has been speculated that VEGF may function as a tumor angiogenesis factor in vivo. Two additional proteins designated VEGF-B and VEGF-C share a significant degree of homology with VEGF. VEGF-B is abundantly expressed in heart and skeletal muscle and is frequently co-expressed with VEGF. VEGF-C binds to and specifically activates Flt-4 and Flk-1. The genes that encode VEGF-B and VEGF-C have been localized to chromosomes 11q13 and 4q34, respectively.

**Synonyms:**

VEGFB, VRF, Vascular endothelial growth factor B, VEGF-related factor

**Product images:**

Western blot (WB) analysis of VEGFB antibody in extracts from Jurkat cells.