

Product datasheet for AP06495PU-N

A RAF (ARAF) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

IHC, WB **Applications:**

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Synthetic peptide, corresponding to amino acids 261-310 of Human Raf. Immunogen:

This antibody detects endogenous levels of A-RAF protein. Specificity:

(region surrounding Asp297)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 68 kDa

Gene Name: A-Raf proto-oncogene, serine/threonine kinase

Database Link: Entrez Gene 369 Human

P10398



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:


Several serine/threonine protein kinases have been implicated as intermediates in signal transduction pathways. These include ERK/MAP kinases, ribosomal S6 kinase (Rsk) and Raf-1. Raf-1 is a cytoplasmic protein with intrinsic serine/threonine activity. It is broadly expressed in nearly all cell lines tested to date and is the cellular homolog of v-Raf, the product of the transforming gene of the 3611 strain of murine sarcoma virus. The unregulated kinase activity of the v-Raf protein has been associated with transformation and mitogenesis while the activity of Raf-1 is normally suppressed by a regulatory N-terminal domain. A-Raf, a second member of the Raf gene family of serine/ threonine protein kinases, exhibits substantial homology to Raf-1 within the kinase domain of the two molecules, but less homology elsewhere. Expression of A-Raf is found at highest levels in urogenital tissues and kidney and at lowest level in brain tissue.

Synonyms: A-raf-1, ARAF1, PKS, PKS2

Product images:



Western blot (WB) analysis of A-RAF antibody in extracts from HeLa cells treated with PMA 125ng/ml 30'.



Immunohistochemistry (IHC) analyzes of A-RAF antibody in paraffin-embedded human breast carcinoma tissue.