

Product datasheet for **AP06513PU-N**

DDIT3 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunofluorescence: 1/50-1/200. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids N-terminus of Human CHOP.
Specificity:	This antibody detects endogenous levels of CHOP protein. (region surrounding Gln26)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Predicted Protein Size:	~ 19, 27 kDa
Gene Name:	DNA damage inducible transcript 3
Database Link:	Entrez Gene 13198 Mouse Entrez Gene 1649 Human P35638



[View online »](#)

Background:

GADD153 is a small nuclear protein that is capable of dimerizing with transcription factors C/EBP alpha and beta. Once dimerized, this complex inhibits the normal binding and function of C/EBP to classical binding sites. Inversely, the C/EBP GADD153 dimer gains binding activity to other non classical C/EBP stress related targets. Under normal cellular conditions this protein is not expressed in detectable levels, but is highly unregulated during times of cellular/ER stress. Examples of GADD153 inducing stress include: treatment with tunicamycin, nutrient starvation and reducing agents that interfere with the calcium flux across the ER membrane.

Synonyms:


DDIT3, DDIT-3, CHOP10, CHOP-10

Protein Families:


Druggable Genome, Transcription Factors

Protein Pathways:

MAPK signaling pathway

Product images:


Immunohistochemistry (IHC) analyzes of CHOP antibody in paraffin-embedded human skeletal muscle tissue.



Western blot (WB) analysis of CHOP antibody in extracts from Jurkat cells.