

## Product datasheet for **AP06540PU-M**

### **MST1 (+MST2) Rabbit Polyclonal Antibody**

#### **Product data:**

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	<b>Western blot:</b> 1/500-1/2000. <b>Immunohistochemistry on paraffin sections:</b> 1/50-1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 150-200 of Human MST1.
Specificity:	This antibody detects endogenous levels of MST1/2 protein. (region surrounding Meth178)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 65 kDa
Gene Name:	macrophage stimulating 1
Database Link:	<a href="#">Entrez Gene 15235 Mouse</a> <a href="#">Entrez Gene 4485 Human</a> <a href="#">P26927</a>



[View online »](#)

**Background:**


Sterile-20 (STE20) is a serine/threonine kinase in *Saccharomyces cerevisiae* that is involved in relaying signals from G protein coupled receptors to cytosolic MAP kinase cascades. Mammalian protein kinases that display sequence similarity to STE20 are divided into two groups, the PAK subfamily and the GCK subfamily. The PAK subfamily members contain a C-terminal catalytic domain and an N-terminal regulatory domain with a p21Rac/Cdc42-binding site, and these kinases can activate both p38MAPK and JNK. The GCK subfamily members contain a C-terminal regulatory domain and an N-terminal catalytic domain, and they have diverse roles in many pathways, including the activation of ERK, JNK, p38 MAPK, and caspase-3. The mammalian STE20-like kinases (MST kinases, also known as Ksr proteins) are members of the GCK subfamily.

**Synonyms:**

MST1, D3F15S2, DNF15S2, HGFL, Macrophage-stimulating protein, Hepatocyte growth factor-like protein, MSP

**Protein Families:**

Druggable Genome

**Product images:**


Immunohistochemistry (IHC) analysis of MST1/2 antibody in paraffin-embedded human breast carcinoma tissue.