

Product datasheet for AP06557PU-S

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

PBK Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to the N-terminal of Human PBK.

Specificity: This antibody detects endogenous levels of PBK/TOPK protein.

(region surrounding Ile4)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 36 kDa

Gene Name: PDZ binding kinase

Database Link: Entrez Gene 55872 Human

Q96KB5






Background:

PBK (PDZ binding kinase) belongs to the serine/threonine protein kinase family. It is related to the dual specific mitogen-activated protein kinase kinase (MAPKK) family and phosphorylates MAP kinase p38. It is active only in mitosis. When phosphorylated it forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin-induced DNA damage. It makes a kinase-substrate complex with cdk1/cyclin B1 and PRC1 on microtubules during mitosis, enhances the cdk1/cyclin B1-dependent phosphorylation of PRC1 and thereby strongly promotes cytokinesis. It may play a role in the activation of lymphoid cells.


Synonyms:

CT84, Nori-3, PDZ-binding kinase, SPK

Product images:



Western blot (WB) analysis of PBK/TOPK antibody in extracts from COS7 cells treated with Nocodazole 1ug/ml 16h.



Immunohistochemistry (IHC) analysis of PBK/TOPK antibody in paraffin-embedded human breast tissue.