

Product datasheet for **AP06565PU-N**

PKC zeta (PRKCZ) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. Immunofluorescence: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 520-570 of Human PKC ζ .
Specificity:	This antibody detects endogenous levels of PKC zeta protein. (region surrounding Pro556)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 67, 78 kDa
Gene Name:	protein kinase C zeta
Database Link:	Entrez Gene 5590 Human Q05513




[View online »](#)

Background:

Members of the protein kinase C (PKC) family play a key regulatory role in a variety of cellular functions including cell growth and differentiation, gene expression, hormone secretion and membrane function. PKCs were originally identified as serine/threonine protein kinases whose activity was dependent on calcium and phospholipids. Diacylglycerols (DAG) and tumor promoting phorbol esters bind to and activate PKC. PKCs can be subdivided into at least two major classes including conventional (c) PKC isoforms (α , β I, β II and γ) and novel (n) PKC isoforms (δ , ϵ , ζ , η and θ). Patterns of expression for each PKC isoform differs among tissues and PKC family members exhibit clear differences in their cofactor dependencies. For instance, the kinase activities of nPKC δ and ϵ are independent of Ca^{++} . On the other hand, nPKC δ and ϵ , as well as all of the cPKC members, possess phorbol ester-binding activities and kinase activities.

Synonyms:

nPKC-zeta, PKC-zeta, PKC zeta, PKC2, Protein kinase C zeta type

Product images:

Western blot (WB) analysis of PKC ζ antibody at 1/500 dilution. Lane 1: HeLa whole cell lysate Lane 2: NIH-3T3 whole cell lysate Lane 3: PC12 whole cell lysate Lane 4: Rat kidney tissue lysate Lane 5: Rat heart tissue lysate Lane 6: PKC ζ



Immunohistochemistry (IHC) analyzes of PKC ζ antibody in paraffin-embedded human brain tissue.