

Product datasheet for **AP06596PU-S**

ZAP70 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western blot: 1/500-1/1000.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 250-300 of Human ZAP70.
Specificity:	This antibody detects endogenous levels of ZAP-70 protein. (region surrounding Thr286)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 70 kDa
Gene Name:	zeta chain of T cell receptor associated protein kinase 70kDa
Database Link:	Entrez Gene 7535 Human P43403




[View online »](#)

Background:

The activation of T lymphocytes by antigens is mediated by the T cell receptor (TCR) which is a multisubunit complex assembled from at least six different genes. The TCR subunits include the Ti α and β chains, the CD3 γ , δ and ϵ chains and a ζ -containing homodimer or heterodimer. The disulfide-linked Ti α - β heterodimer is responsible for antigen recognition, but the short five amino acid cytoplasmic domains of Ti α and β are unlikely to be sufficient to couple to intracellular signaling pathways. In contrast, the structured features of the CD3 and ζ subunits suggest a role in signal transduction. Of these, the ζ chain, which is expressed as either a homodimer or heterodimer, has a short extracellular domain of only 9 amino acids, but a larger 113 amino acid cytoplasmic domain. A 70 kDa tyrosine phosphoprotein, ZAP-70, has been identified that associates with ζ and undergoes tyrosine phosphorylation following TCR stimulation.

Synonyms:

ZAP-70, ZAP 70, SRK

Product images:

Western blot (WB) analysis of ZAP-70 antibody in extracts from 293 cells treated with PMA 125ng/ml 30'.