

Product datasheet for **AP06600PU-S**

FAK (PTK2) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 370-420 of Human FAK.
Specificity:	This antibody detects endogenous levels of FAK protein. (region surrounding Ile401)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 15 mM sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 125 kDa
Gene Name:	protein tyrosine kinase 2
Database Link:	Entrez Gene 5747 Human Q05397




[View online »](#)

Background:


Focal adhesion kinase was initially identified as a major 125 kDa substrate for the intrinsic protein tyrosine kinase activity of Src-encoded pp60. The deduced amino acid sequence of FAK p125 has shown it to be a cytoplasmic protein tyrosine kinase whose sequence and structural organization are unique as compared to other proteins described to date. Localization of p125 by immunofluorescence suggests that it is primarily found in cellular focal adhesions leading to its designation as focal adhesion kinase (FAK). FAK is concentrated at the basal edge of only basal keratinocytes that are actively migrating and rapidly proliferating in repairing burn wounds, and is activated and localized to the focal adhesions of spreading keratinocytes in culture. Thus, it has been postulated that FAK may have an important in vivo role in the re-epithelialization of human wounds. FAK protein tyrosine kinase activity has also been shown to increase in cells stimulated to grow by use of mitogenic neuropeptides or neurotransmitters acting through G protein-coupled receptors.

Synonyms:

FAK, Focal adhesion kinase 1, FADK1, pp125FAK, Protein-tyrosine kinase 2

Product images:

Western blot (WB) analysis of FAK antibody in extracts from hela cells.



Immunohistochemistry (IHC) analyzes of FAK antibody in paraffin-embedded human brain tissue.