

Product datasheet for AP06619PU-M

MAPKAP Kinase 2 (MAPKAPK2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies IHC, WB **Applications:** Recommended Dilution: Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. **Reactivity:** Human, Mouse, Rat Host: Rabbit **Clonality:** Polyclonal Synthetic peptide, corresponding to amino acids 300-350 of Human MAPKAPK-2. Immunogen: This antibody detects endogenous levels of MAPKAPK-2 protein. Specificity: (region surrounding Ser328) Formulation: Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified lg fraction Preservative: 15 mM sodium azide **Concentration:** 1.0 mg/ml **Purification:** Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE) **Conjugation:** Unconjugated Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Stability: Shelf life: one year from despatch. **Predicted Protein Size:** ~ 46 kDa Gene Name: mitogen-activated protein kinase-activated protein kinase 2 Database Link: Entrez Gene 9261 Human P49137



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	MAPKAP Kinase 2 (MAPKAPK2) Rabbit Polyclonal Antibody – AP06619PU-M
Background:	MAP kinase activated protein kinase 2 (MAPKAP Kinase 2), also known as p45 hsp27 kinase, is a 45-54 kDa serine/threonine protein kinase that contains a proline rich sequence and two putative SH3 binding sites. MAPKAP Kinase 2 is activated in response to stress, IL1 and TNF, possibly catalyzed by p38/Hog dependent phosphorylation. One of the major substrates of MAPKAP Kinase 2 is hsp27, which stimulates actin polymerization in order to facilitate recovery from destruction of cytoskeleton during cellular stresses. MAPKAP2 is implicated in several disorders including ischemic brain injury and heart failure and has been shown to be important in regulating stress resistance and the production of TNF alpha.


Synonyms:

MAPK-activated protein kinase 2, MAPKAP kinase 2, MAPKAPK-2, MK2, MAPKAPK2

Product images:



Western blot (WB) analysis of MAPKAPK-2 antibody in extracts from raw264.7 cells.



Immunohistochemistry (IHC)analyzesof MAPKAPK-2 antibody in paraffin-embedded human breast carcinoma tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US