

Product datasheet for AP06621PU-N

BID Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

IHC, WB **Applications:**

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on Paraffin Sections: 1/50-1/200.

Reactivity: Human Host: Rabbit

Clonality: Polyclonal

Synthetic peptide, corresponding to amino acids 50-100 of Human BID. Immunogen:

This antibody detects endogenous levels of BID protein. Specificity:

(region surrounding Ile72)

Formulation: Phosphate buffered saline (PBS), pH~7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)

Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage:

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~22 kDa

Gene Name: BH3 interacting domain death agonist

Database Link: Entrez Gene 637 Human

P55957



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:


Bid, a BH3 domain containing proapoptotic Bcl2 family member, is localized in the cytosolic fraction of cells as an inactive precursor. Its active form is generated upon proteolytic cleavage by caspase 8 in the Fas signaling pathway. Cleaved Bid translocates to mitochondria and releases its potent proapoptotic activity, which in turn induces cytochrome c release and mitochondrial damage. The cytochrome c releasing activity of Bid was antagonized by Bcl2. Mutation in the SH3 domain can diminish the cytochrome c releasing activity. In animal model studies, Bid deficient mice are found resistant to the lethal effects of death factor signals relayed through Fas.

Synonyms: p22 BID

Product images:



Western blot (WB) analysis of BID antibody at 1/500 dilution Lane 1:MCF-7 cell lysate Lane 2:NIH-3T3 cell lysate Lane 3:H9C2 cell lysate



Immunohistochemistry analysis of BID Antibody in paraffin-embedded human brain tissue.