

## Product datasheet for **AP06630PU-S**

### Thrombin Receptor (F2R) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	<b>Western blot:</b> 1/500-1/1000.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to the N-terminal of Human Thrombin R.
Specificity:	This antibody detects endogenous levels of PAR1 protein. (region surrounding Gly17)
Formulation:	Phosphate buffered saline (PBS), pH 7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 48 kDa
Gene Name:	coagulation factor II thrombin receptor
Database Link:	<a href="#">Entrez Gene 2149 Human P25116</a>




[View online »](#)

**Background:**

Thrombin receptor (also designated protease-activated receptor-1 or PAR-1), PAR-2 and PAR-3 compose a distinct class of G protein-coupled receptors activated by proteolysis. Cleavage of these receptors by proteases occurs within the amino-terminal extracellular domain. Thrombin, a serine protease involved in platelet aggregation and blood coagulation, activates the thrombin receptor, resulting in elevated intracellular calcium levels in platelets. Thrombin also cleaves PAR-3 in vitro, suggesting that PAR-3 may be involved in thrombosis or mitogenesis. Thrombin receptor and PAR-4 appear to account for most thrombin signaling in platelets. Activation of PAR-2 in vitro is induced by trypsin, suggesting that PAR-2 is not an alternative thrombin receptor. Cytokines including TNF-alpha and IL-1beta increase PAR-2 expression, indicating PAR-2 involvement in the acute inflammatory response.

**Synonyms:**

Proteinase-activated receptor 1, PAR-1, PAR1, CF2R

**Product images:**

Western blot (WB) analysis of PAR1 antibody in extracts from HeLa cells treated with Nocodazole 1ug/ml 18 hours.