

Product datasheet for AP06638PU-N

AIRE Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IF, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunofluorescence: 1/50-1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 101-150 of Human AIRE.
Specificity:	This antibody detects endogenous levels of AIRE-1 protein. (region surrounding Pro126)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS- PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 60 kDa
Gene Name:	autoimmune regulator
Database Link:	<u>Entrez Gene 326 Human</u> <u>O43918</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE AIRE Rabbit Polyclonal Antibody – AP06638PU-N

Background: The autoimmune regulator gene, which is defective in the hereditary autoimmune disease APECED, encodes the transcriptional activator AIRE. AIRE is expressed in the medullary epithelial cells and monocyte-dendritic cells of the thymus, with lower expression in the spleen, fetal liver and lymph nodes. In adult tissue, AIRE expression in the thymus is confined to the medulla and the cortico-medullary junction, where it is modulated by thymocytes undergoing negative selection. At the cellular level, AIRE is located in microtubular structures of the cytoskeleton and in discrete nuclear dots resembling ND10 nuclear bodies. AIRE is induced by developing early thymocytes and is associated with the correct establishment of a regular thymic environment. AIRE regulates thymic architecture via transcriptional control of downstream target genes. AIRE mutations in APECED patients may affect thymic T cell selection and the formation of self-tolerance.

Synonyms:

Autoimmune regulator

Product images:



Western blot (WB) analysis of AIRE antibody at 1/500 dilution Lane 1: Jurkat whole cell lysate Lane 2: Mouse lung tissue lysate Lane 3: Rat lung tissue lysate

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US