

Product datasheet for **AP06657PU-N**

p53 (TP53) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. Immunofluorescence: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 301-350 of Human p53.
Specificity:	This antibody detects endogenous levels of p53 protein. (region surrounding Leu330)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 53, 43 kDa
Gene Name:	tumor protein p53
Database Link:	Entrez Gene 7157 Human P04637



[View online »](#)


Background:

p53, a DNA-binding, oligomerization domain and transcription activation domain-containing tumor suppressor, upregulates growth arrest and apoptosis-related genes in response to stress signals, thereby influencing programmed cell death, cell differentiation and cell cycle control mechanisms. p53 localizes to the nucleus, yet can be chaperoned to the cytoplasm by the negative regulator MDM2, an E3 ubiquitin ligase that is upregulated in the presence of active p53, where MDM2 poly-ubiquitinates p53 for proteasome targeting. p53 fluctuates between latent and active (DNA-binding) conformations and is differentially activated through posttranslational modifications including phosphorylation and acetylation. Mutations in the DNA-binding domain (DBD, amino acids 110-286) of p53 can compromise energetically favorable association with cis elements and are implicated in several human cancers.


Synonyms:

Cellular tumor antigen p53, Tumor suppressor p53, Phosphoprotein p53, NY-CO-13

Product images:



Western blot (WB) analysis of p53 antibody in extracts from HepG2 cells.



Immunohistochemistry (IHC) analysis of p53 antibody in paraffin-embedded human brain tissue.