

Product datasheet for AP06658PU-M

Creatine kinase M type (CKM) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies Applications: IF, IHC, WB Recommended Dilution: Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. Immunofluorescence: 1/50-1/200. **Reactivity:** Human, Mouse, Rat Host: Rabbit **Clonality:** Polyclonal Immunogen: Synthetic peptide, corresponding to the N-terminus of Human Creatine Kinase M. Specificity: This antibody detects endogenous levels of Creatine Kinase M protein. (region surrounding Tyr14) Formulation: Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide **Concentration:** 1.0 mg/ml **Purification:** Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE) **Conjugation:** Unconjugated Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Storage: Avoid repeated freezing and thawing. Stability: Shelf life: one year from despatch. **Predicted Protein Size:** ~ 43 kDa Gene Name: creatine kinase, M-type Database Link: Entrez Gene 12715 MouseEntrez Gene 24265 RatEntrez Gene 1158 Human P06732



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Screatine kinase M type (CKM) Rabbit Polyclonal Antibody – AP06658PU-M

Background:	Creatine kinase isoenzymes play a central role in energy transduction in tissues with large,
	fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa. The CKM
	isoform, predominant in skeletal muscle and heart tissue, is a cytoplasmic enzyme involved
	in energy homeostasis and is an important serum marker for myocardial infarction. CKM
	reversibly catalyzes the transfer of phosphate between ATP and various phosphogens such
	as creatine phosphate. It acts as a homodimer in striated muscle as well as in other tissues,
	and as a heterodimer with a similar brain isozyme in heart. The encoded protein is a member
	of the ATP:guanido phosphotransferase protein family.

Synonyms:Creatine kinase M-type, Creatine kinase M chain, M-CK, CKM, Creatine kinase MMProtein Families:Druggable GenomeProtein Pathways:Arginine and proline metabolism, Metabolic pathways

Product images:



Western blot (WB) analysis of Creatine Kinase M antibody at 1/500 dilution Lane 1:Hela whole cell lysate Lane 2:NIH-3T3 whole cell lysate Lane 3:H9C2 whole cell lysate



Immunohistochemistry (IHC) analyzes of Creatine Kinase M antibody in paraffin-embedded human heart tissue.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US