

## Product datasheet for **AP06677PU-N**

### PFKL Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	<b>Western blot:</b> 1/500-1/1000. <b>Immunohistochemistry on Paraffin Sections:</b> 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 700-750 of Human PFKL.
Specificity:	This antibody detects endogenous levels of PFK-B protein. (region surrounding Glu724)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~85 kDa
Gene Name:	phosphofructokinase, liver type
Database Link:	<a href="#">Entrez Gene 18641 Mouse</a> <a href="#">Entrez Gene 25741 Rat</a> <a href="#">Entrez Gene 5211 Human</a> <a href="#">P17858</a>



[View online »](#)

**Background:**

Phosphofructokinases (PFK) are regulatory glycolytic enzymes that convert fructose 6-phosphate and ATP into fructose 1,6-bisphosphate (through PFK-1), fructose 2,6-bisphosphate (through PFK-2) and ADP. Human PFK-1 is tetrameric and isoenzymes include PFK-1 muscle (PFKM, PFK-A), PFK-1 liver (PFKL, PFK-B) and PFK-1 platelet (PFKP, PFK-C, PFKF). PFK-1 is inhibited by ATP and citrate (from the tricarboxylic acid cycle). PFK-1 undergoes activation in the presence of elevated AMP. The most potent activator is fructose 2,6-bisphosphate, which is produced by PFK-2 from the same substrate, fructose 6-phosphate. PFK-2 is bifunctional and a key regulator for PFK-1. PFK-2 catalyzes the synthesis of fructose 2,6-bisphosphate and contains fructose 2,6-bisphosphatase activity that catalyzes the degradation of fructose 2,6-bisphosphate. PFK-2 is dimeric and isoenzymes include PFK-2 liver (PFKFB1, PFRX), PFK-2 cardiac (PFKFB2), PFK-2 placental (PFKFB3, inducible PFK-2) and PFK-2 testis (PFKFB4).

**Synonyms:**


Phosphofructokinase 1, 6-phosphofructokinase liver, PFK-B, Phosphohexokinase

**Protein Families:**


Druggable Genome

**Protein Pathways:**

Fructose and mannose metabolism, Galactose metabolism, Glycolysis / Gluconeogenesis, Metabolic pathways, Pentose phosphate pathway

**Product images:**


Western blot analysis of PFK-B Antibody in extracts from A549 cells.



Immunohistochemistry analyzes of PFK-B  
Antibody in paraffin-embedded human brain  
tissue.