

Product datasheet for AP06681PU-M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cleavage stimulation factor 2 (CSTF2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immunofluorescence: 1/50-1/200.

Reactivity: Human, Mouse

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids N-terminus of Human CstF-64.

Specificity: This antibody detects endogenous levels of CstF-64 protein.

(region surrounding Arg46)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 64 kDa

Gene Name: cleavage stimulation factor subunit 2

Database Link: Entrez Gene 108062 MouseEntrez Gene 1478 Human

P33240



Cleavage stimulation factor 2 (CSTF2) Rabbit Polyclonal Antibody - AP06681PU-M


Background:

The serine threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet derived growth factor. The activation is rapid and specific. In the developing nervous system AKT is a critical mediator of growth factor induced neuronal survival. Survival factors can suppress apoptosis in a transcription independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Multiple alternatively spliced transcript variants have been found for this gene (referenced from entrez gene).

Synonyms:

CSTF 64 kDa subunit, CstF-64, CSTF64, CF-1 64 kDa subunit

Product images:



Western blot (WB) analysis of CstF-64 antibody in extracts from 293 cells.