

Product datasheet for AP06682PU-S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRK5 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 350-400 of Human GRK5.

Specificity: This antibody detects endogenous levels of GRK5 protein.

(region surrouding Gln382)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~68 kDa

Gene Name: G protein-coupled receptor kinase 5

Database Link: Entrez Gene 2869 Human

P34947






Background:

Heterotrimeric G protein-mediated signal transduction is a dynamically regulated process with the intensity of signal decreasing over time despite the continued presence of the agonist. This phenomenon, referred to as agonistmediated desensitization, involves phosphorylation of the receptor by two classes of enzymes. The first are the second messenger-regulated kinases such as c-AMP dependent protein kinase A and protein kinase C. The second are the G protein-coupled receptor kinases (GRKs). At least seven members of the GRK family have been identified. These include rhodopsin kinase, GRK 1; two forms of β -adrenergic receptor kinase, GRK 2 (β ARK, β ARK1) and GRK 3 (β ARK2); IT-11 (GRK 4); GRK 5, GRK 6 and GRK 7. Phosphorylation of receptors by GRKs appears to be strictly dependent on the receptor being in its agonist-activated state.

Synonyms:

G protein-coupled receptor kinase 5, GRK5

Product images:



Western blot (WB) analysis of GRK5 antibody in extracts from HeLa cells.