

Product datasheet for **AP06689PU-M**

HNF 4 alpha (HNF4A) (+ gamma) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 100-150 of Human HNF4α.
Specificity:	This antibody detects endogenous levels of HNF-4α/γ protein. (region surrounding Lys127)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 52 kDa
Gene Name:	hepatocyte nuclear factor 4 alpha
Database Link:	Entrez Gene 15378 Mouse Entrez Gene 25735 Rat Entrez Gene 3172 Human P41235



[View online »](#)

Background:

HNF-1 (α and β), HNF-3 (α , β and γ), HNF-4 (α and γ), and HNF-6 compose, in part, a homoeoprotein family designated the Hepatocyte Nuclear Factor family. The various HNF-1 isoforms regulate transcription of genes in the liver as well as in other tissues such as kidney, small intestine and thymus. HNF-3 α , HNF-3 β and HNF-3 γ regulate the transcription of numerous hepatocyte genes in adult liver. HNF-3 α and HNF-3 β have also been shown to be involved in gastrulation events such as body axis formation. HNF-4 α and HNF-4 γ have been shown to be important for early embryo development. HNF-4 α is expressed in liver, kidney, pancreas, small intestine, testis and colon; and HNF-4 γ is expressed in each of these tissues except liver. HNF-6 has been shown to bind to the promoter of HNF-3 β , which indicates a potential role of HNF-6 in gut endoderm epithelial cell differentiation. Evidence suggests that HNF-6 may also be a transcriptional activator for at least 22 other hepatocyte-enriched genes, including cytochrome P450 2C13 and α -1 antitrypsin.

Synonyms:

HNF4A, HNF4, NR2A1, Hepatocyte nuclear factor 4-alpha, Transcription factor HNF-4, Transcription factor 14


Protein Families:

Druggable Genome, ES Cell Differentiation/IPS, Nuclear Hormone Receptor, Transcription Factors


Protein Pathways:

Maturity onset diabetes of the young

Product images:



Western blot (WB) analysis of HNF-4a/? antibody in extracts from A549 cells.



Immunohistochemistry (IHC) analysis of HNF-4a/? antibody on paraffin-embedded sections.