

Product datasheet for AP06692PU-M

GRK6 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IF, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunofluorescence: 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids N-terminus of Human GRK 6.
Specificity:	This antibody detects endogenous levels of GRK 6 protein. (region surrounding asn22)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS- PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 65 kDa
Gene Name:	G protein-coupled receptor kinase 6
Database Link:	<u>Entrez Gene 2870 Human</u> <u>P43250</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRK6 Rabbit Polyclonal Antibody – AP06692PU-M

Background:Heterotrimeric G protein-mediated signal transduction is a dynamically regulated process
with the intensity of signal decreasing over time despite the continued presence of the
agonist. This phenomenon, referred to as agonistmediated desensitization, involves
phosphorylation of the receptor by two classes of enzymes. The first are the second
messenger-regulated kinases such as c-AMP dependent protein kinase A and protein kinase
C. The second are the G protein-coupled receptor kinases (GRKs). At least seven members of
the GRK family have been identified. These include rhodopsin kinase, GRK 1; two forms of β-
adrenergic receptor kinase, GRK 2 (βARK, βARK1) and GRK 3 (βARK2); IT-11 (GRK 4); GRK 5,
GRK 6 and GRK 7. Phosphorylation of receptors by GRKs appears to be strictly dependent on
the receptor being in its agonist-activated state.

Synonyms: G protein-coupled receptor kinase 6, GPRK6

Product images:



Western blot (WB) analysis of GRK 6 antibody in extracts from Jurkat cells.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US