

Product datasheet for AP06695PU-M

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

YAP1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunofluorescence: 1/50-1/200.

Immunohistochemistry on Paraffin Sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 280-330 of Human YAP.

Specificity: This antibody detects endogenous levels of YAP protein.

(region surrounding Gln295)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography using epitope-specific immunogen

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~38, 67 kDa

Gene Name: Yes associated protein 1

Database Link: Entrez Gene 10413 Human

P46937




Background:


The Yes-associated protein, otherwise known as YAP, is a 14-3-3-binding molecule that was originally recognized by virtue of its ability to bind to the SH3 domain of Yes. The binding of YAP to 14-3-3 requires the phosphorylation of a homologous serine residue (Ser 112) in the YAP 14-3-3-binding motif. The highly conserved and ubiquitously expressed 14-3-3 proteins regulate differentiation, cell cycle progression and apoptosis by binding intracellular phosphoproteins involved in signal transduction. YAP may link events at the plasma membrane and cytoskeleton to inhibition of transcription in the nucleus in a manner regulated by 14-3-3 proteins. YAP shares homology with the WW domain of TAZ, transcriptional co-activator with PDZ-binding motif, which functions as a transcriptional co-activator by binding to the PPXY motif present in transcription factors. YAP is expressed at high levels in the lung, placenta, prostate, ovary and testis.

Synonyms: YAP, YAP65

Product images:



Western blot analysis of YAP Antibody in extracts from HeLa cells.



Immunohistochemistry analysis of YAP Antibody in paraffin-embedded human breast carcinoma tissue.