

Product datasheet for AP06700PU-M

MEOX 2 (MEOX2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse, Rat

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 150-200 of Human MOX-2.

Specificity: This antibody detects endogenous levels of MOX-2 protein.

(region surrounding asn184)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 33 kDa

Gene Name: mesenchyme homeobox 2

Database Link: Entrez Gene 4223 Human

P50222



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Closely related homeobox proteins, MOX-1 and MOX-2, belong to a family of nonclustered, diverged homeobox genes that are expressed in overlapping patterns in the paraxial mesoderm and its derivatives. MOX-1 and MOX-2 function transiently in the formation of mesodermal and mesenchymal derivatives. MOX-1 and MOX-2 are implicated in the early steps of mesoderm formation during gastrulation. In addition, the MOX proteins are also involved in somatic differentiation. Significantly, MOX-1 associates more strongly with Pax1, whereas MOX-2 preferentially associates with Pax3. Specifically, expression of MOX-2, also known as Mesenchyme homeobox 2 and GAX, has been shown to be critical in axial skeleton development. MOX-2 is not needed for the migration of myogenic precursors into the limb bud, but it is essential for normal appendicular muscle formation and for the normal regulation of myogenic genes. MOX-2 is expressed in placental tissue. The human MEOX2 gene maps to chromosome 7p22.1-p21.3 and encodes the MOX-2 protein. Mutations in the gene may be involved in craniofacial and/or skeletal abnormalities.

Synonyms:

GAX, MOX2, Growth arrest-specific homeobox, Mesenchyme homeobox 2, Homeobox protein MOX-2

Product images:



Western blot (WB) analysis of MOX-2 antibody in extracts from COLO205 cells.