

## Product datasheet for AP06705PU-M

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## MEK6 (MAP2K6) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IF, IHC, WB

Recommended Dilution: Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Immuofluorescence: 1/50-1/200.

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

Clonality: Polyclonal

**Immunogen:** Synthetic peptide, corresponding to amino acids N-terminus of Human MEK-6.

**Specificity:** This antibody detects endogenous levels of MEK-6 protein.

(region surrounding Gln3)

**Formulation:** Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified lg fraction Preservative: 0.05% sodium azide

Concentration: 1.0 mg/ml

**Purification:** Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-

PAGE)

Conjugation: Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: ~ 37 kDa

**Gene Name:** mitogen-activated protein kinase kinase 6

**Database Link:** Entrez Gene 5608 Human

P52564






#### Background:

A family of protein kinases located upstream of the MAP kinases and responsible for their activation has been identified. The prototype member of this family, designated MAP kinase kinase, or MEK-1, specifically phosphorylates the MAP kinase regulatory threonine and tyrosine residues present in the Thr-Glu-Tyr motif of ERK. A second MEK family member, MEK-2, resembles MEK-1 in its substrate specificity. MEK-3 (or MKK-3) functions to activate p38 MAP kinase, and MEK-4 (also called SEK1 or MKK-4) activates both p38 and JNK MAP kinases. MEK-5 appears to specifically phosphorylate ERK5, whereas MEK-6 phosphorylates p38 and p38b. MEK-7 (or MKK-7) phosphorylates and activates the JNK signal transduction pathway.

Synonyms:

MAPKK 6, MAPK/ERK kinase 6, SAPKK3, MEK6, MKK6, MAP kinase kinase 6

# **Product images:**



Western blot (WB) analysis of MEK-6 antibody in extracts from 293 cells.