

## Product datasheet for **AP06719PU-N**

### NCBP1 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	<b>Western blot:</b> 1/500-1/1000. <b>Immunohistochemistry on paraffin sections:</b> 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to the N-terminus of Human CBP80.
Specificity:	This antibody detects endogenous levels of CBP80 protein. (region surrounding Arg5)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 15 mM sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 88 kDa
Gene Name:	nuclear cap binding protein subunit 1
Database Link:	<a href="#">Entrez Gene 4686 Human Q09161</a>


[View online »](#)

**Background:**


In eukaryotes, the majority of mRNAs have an m(7)G cap, which is added cotranscriptionally and plays a critical role in many aspects of mRNA metabolism. The effect of the cap on translation is mediated by the initiation factor eIF-4E, whereas the effect on pre-mRNA splicing involves a nuclear complex(CBC). CBC consists of two cap binding proteins CBP20 and CBP80, which mediate the stimulatory functions of the cap in pre-mRNA splicing, 3' end formation and U snRNA export. The genes CBC1 and CBC2 encode CBP80 and CBP20, respectively. CBP80 comprises three domains, each containing a MIF4G domain. CBP20 has an RNAP fold and associates with the second and third domains of CBP80. CBP also plays a role in nonsense-mediated decay(NMD), which eliminates mRNAs, which prematurely terminate translation.CBP80-bound mRNA undergoes a “pioneer” round of translation beforeCBP80-CBP20 are replaced by eIF4E, and Upf2 and Upf3 proteins.

**Synonyms:**

CBP80, NCBP

**Product images:**


Western blot (WB) analysis of CBP80 antibody at 1/500 dilution Lane 1:MCF-7 whole cell lysate  
Lane 2:Raw264.7 whole cell lysate Lane 3:PC12 whole cell lysate



Immunohistochemistry (IHC) analyzes of CBP80 antibody in paraffin-embedded colon carcinoma tissue.