

Product datasheet for **AP06743PU-M**

PIG3 (TP53I3) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western blot: 1/500-1/1000.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 230-286 of Human PIG3.
Specificity:	This antibody detects endogenous levels of PIG3 protein. (region surrounding Leu265)
Formulation:	Phosphate buffered saline (PBS), pH 7.2. State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 36 kDa
Gene Name:	tumor protein p53 inducible protein 3
Database Link:	Entrez Gene 9540 Human Q53FA7




[View online »](#)

Background:

The PIG (p53-induced gene) gene family encodes redox-controlling proteins that are involved in p53 tumor suppressor activity. One member of the PIG gene family, p53-inducible gene 3 (PIG3), is a p53 responsive gene that maps, in humans, to chromosome 2p and encodes a protein with significant homology to oxidoreductases. Oxidoreductases are enzymes involved in cellular responses to oxidative stress and irradiation, and they influence the involvement of PIG3 in the metabolism of reactive oxygen species. PIG3 is localized to the cytoplasm and induced in primary, non-transformed, and transformed cell cultures after exposure to genotoxic agents. The induction of PIG3 is p53 dependent and occurs with delayed kinetics as compared with other p53 downstream targets. PIG3 may act with caspase-8 as a key regulatory element in p53-dependent transcriptional deregulation by triggering the caspase cascade and mitochondrial breakdown. PIG3 is highly up-regulated by p53 and may be useful for detecting transient activation of p53.

Synonyms:

p53-induced gene 3 protein, PIG3

Product images:


Western blot (WB) analysis of PIG3 antibody in extracts from 293 cells.