

Product datasheet for **AP06748PU-S**

EZH1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunofluorescence: 1/50-1/200. Immunohistochemistry on Paraffin Sections: 1/50-1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 162-208 of Human ENX-2.
Specificity:	This antibody detects endogenous levels of ENX-2 protein. (region surrounding Glu192)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~85 kDa
Gene Name:	enhancer of zeste 1 polycomb repressive complex 2 subunit
Database Link:	Entrez Gene 14055 Mouse Entrez Gene 2145 Human Q92800



[View online »](#)

Background:


In *Drosophila*, the Polycomb (PcG) gene family encodes chromatin proteins that are required for the repression of homeotic loci in embryonic development. PcG proteins work in conjunction with the trithorax-group (trxG) proteins, which activate homeobox gene expression during embryonic development. ENX-1, a mammalian homolog of the *Drosophila* gene enhancer of zeste, is a PcG protein that is ubiquitously expressed during early embryogenesis and becomes restricted to the central and peripheral nervous systems and sites of fetal hematopoiesis during later development. In the adult, ENX-1 is restricted to specific sites, including spleen, testis and placenta. ENX-2 is another mammalian homolog of the *Drosophila* gene enhancer of zeste and contains one SET domain. The gene for human ENX-2 maps to chromosome 17q21. ENX-2 expression is ubiquitous in adult and fetal tissue, where it may aid in maintaining heterochromatin stability.

Synonyms:


ENX-2, KIAA0388

Protein Families:

Druggable Genome, Transcription Factors

Product images:

Western blot (WB) analysis of ENX-2 antibody in extracts from HT-29 cells.



Immunohistochemistry (IHC) analyzes of ENX-2 antibody in paraffin-embedded human brain tissue.